arXiv:0801.0219v1 [math.FA] 31 Dec 2007

CHARACTERIZATIONS OF ALGEBRAS OF RAPIDLY DECREASING GENERALIZED FUNCTIONS

C. BOUZAR AND T. SAIDI

Abstract. The well-known characterizations of Schwartz space $\mathcal{S}$ of rapidly decreasing functions is extended to the algebra $\mathcal{G}_{\mathcal{S}}$ of rapidly decreasing generalized functions and to the algebra $\mathcal{G}_{\mathcal{S}}^{\infty}$ of regular rapidly decreasing generalized functions.

1. Introduction

The Schwartz space $\mathcal{S}$ of rapidly decreasing functions on $\mathbb{R}^n$ and its generalizations, in view of their importance in many domains of analysis, have been characterized differently by many authors, e.g. see [10], [15], [17], [4], [1] and [11]. To built a Fourier analysis within the generalized functions of [5], the algebra of rapidly decreasing generalized functions on $\mathbb{R}^n$, denoted $\mathcal{G}_{\mathcal{S}}$, was first constructed in [18] and recently studied in [8] and [7]. The algebra of regular rapidly decreasing generalized functions on $\mathbb{R}^n$, denoted $\mathcal{G}_{\mathcal{S}}^{\infty}$, is fundamental in the characterization of the local regularity of a Colombeau generalized function by its Fourier transform and also for developing a generalized microlocal analysis.

Let

$$\mathcal{S}^{*} = \left\{ f \in \mathcal{C}^{\infty} : \forall \alpha \in \mathbb{Z}_{+}^{n}, \sup_{x \in \mathbb{R}^n} \left| \partial^{\alpha} f(x) \right| < \infty \right\},$$

$$\mathcal{S}_{*} = \left\{ f \in \mathcal{C}^{\infty} : \forall \beta \in \mathbb{Z}_{+}^{n}, \sup_{x \in \mathbb{R}^n} \left| x^{\beta} f(x) \right| < \infty \right\},$$

then, inspired by the work of [15], the authors of [4] proved the following result :

$$\mathcal{S} = \mathcal{S}^{*} \cap \mathcal{S}_{*}$$

The aim of this work is to characterize the algebras $\mathcal{G}_{\mathcal{S}}$ and $\mathcal{G}_{\mathcal{S}}^{\infty}$ in the spirit of the characterization of the Schwartz space $\mathcal{S}$ done in [4]. In fact we do more, this characterization is given in the general context of the algebra $\mathcal{G}_{\mathcal{S}}^{\mathcal{R}}(\Omega)$ of $\mathcal{R}-$regular rapidly decreasing generalized functions on an open set Ω of $\mathbb{R}^n$, see [7] and [3]. The sixth section of this paper gives such an extension, i.e. the characterization of the algebra $\mathcal{G}_{\mathcal{S}}^{\mathcal{R}}(\Omega)$ provided Ω is a box of $\mathbb{R}^n$. The seventh section gives a characterization of $\mathcal{G}_{\mathcal{S}}^{\mathcal{R}}(\mathbb{R}^n)$ of $\mathcal{R}-$regular rapidly decreasing generalized functions on the whole space $\mathbb{R}^n$ using the Fourier transform. The last section, as corollaries of the results of the paper, gives the characterizations of the classical algebras $\mathcal{G}_{\mathcal{S}}$ and $\mathcal{G}_{\mathcal{S}}^{\infty}$.

1991 *Mathematics Subject Classification.* 46F30; 46F05; 42B10.

Key words and phrases. Schwartz space S, Rapidly decreasing generalized functions, Colombeau generalized functions, Fourier transform.

2. Regular sets of sequences

We will adopt the notations and definitions of distributions and Colombeau generalized functions, see [14] and [13] .

Definition 1. *A non void subset* $\mathcal{R}$ *of* $\mathbb{R}_+^{\mathbb{Z}_+}$ *is said to be regular, if*

For all $(N_m)_{m\in\mathbb{Z}_+}\in\mathcal{R}$ *and* $(k,k')\in\mathbb{Z}_+^2$*, there exists* $(N'_m)_{m\in\mathbb{Z}_+}\in\mathcal{R}$ *such that*

$$N_{m+k}+k'\leq N'_m\ ,\ \forall m\in\mathbb{Z}_+ \tag{R1}$$

For all $(N_m)_{m\in\mathbb{Z}_+}$ *and* $(N'_m)_{m\in\mathbb{Z}_+}$ *in* $\mathcal{R}$*, there exists* $(N"_m)_{m\in\mathbb{Z}_+}\in\mathcal{R}$ *such that*

$$\max\left(N_m,N'_m\right)\leq N"_m\ ,\ \forall m\in\mathbb{Z}_+ \tag{R2}$$

For all $(N_m)_{m\in\mathbb{Z}_+}$ *and* $(N'_m)_{m\in\mathbb{Z}_+}$ *in* $\mathcal{R}$*, there exists* $(N"_m)_{m\in\mathbb{Z}_+}\in\mathcal{R}$ *such that*

$$N_{l_1}+N'_{l_2}\leq N"_{l_1+l_2}\ ,\ \forall\left(l_1,l_2\right)\in\mathbb{Z}_+^2 \tag{R3}$$

Example 1. *The set* $\mathbb{R}_+^{\mathbb{Z}_+}$ *of all positive sequences is regular.*

Example 2. *The set* $\mathcal{A}$ *of affine sequences defined by*

$$\mathcal{A}=\left\{(N_m)_{m\in\mathbb{N}}\in\mathbb{R}_+^{\mathbb{Z}_+}:\exists a\geq0,\exists b>0,\forall l\in\mathbb{Z}_+,N_l\leq al+b\right\}$$

is regular.

Example 3. *The set* $\mathcal{B}$ *of all bounded sequences of* $\mathbb{R}_+^{\mathbb{Z}_+}$ *is regular.*

The notion of regular set is extended to the sets of double sequences.

Definition 2. *A non void subset* $\widetilde{\mathcal{R}}$ *of* $\mathbb{R}_+^{\mathbb{Z}_+^2}$ *is said to be regular if*

For all $(N_{q,l})_{(q,l)\in\mathbb{Z}_+^2}\in\widetilde{\mathcal{R}}$ *and* $(k,k',k'')\in\mathbb{Z}_+^3$*, there exists* $\left(N'_{q,l}\right)_{(q,l)\in\mathbb{Z}_+^2}\in\widetilde{\mathcal{R}}$ *such that*

$$N_{q+k,l+k'}+k''\leq N'_{q,l}\ ,\ \forall\left(q,l\right)\in\mathbb{Z}_+^2 \tag{$\widetilde{R}1$}$$

For all $(N_{q,l})_{(q,l)\in\mathbb{Z}_+^2}$ *and* $\left(N'_{q,l}\right)_{(q,l)\in\mathbb{Z}_+^2}$ *in* $\widetilde{\mathcal{R}}$ *, there exists* $(N"_{q,l})_{(q,l)\in\mathbb{Z}_+^2}\in\widetilde{\mathcal{R}}$ *such that*

$$\max\left(N_{q,l},N'_{q,l}\right)\leq N"_{q,l}\ ,\ \forall\left(q,l\right)\in\mathbb{Z}_+^2 \tag{$\widetilde{R}2$}$$

For all $(N_{q,l})_{(q,l)\in\mathbb{Z}_+^2}$ *and* $\left(N'_{q,l}\right)_{(q,l)\in\mathbb{Z}_+^2}$ *in* $\widetilde{\mathcal{R}}$ *, there exists* $(N"_{q,l})_{(q,l)\in\mathbb{Z}_+^2}\in\widetilde{\mathcal{R}}$ *such that*

$$N_{q_1,l_1}+N'_{q_2,l_2}\leq N"_{q1+q2,l_1+l_2}\ ,\ \forall\left(q_1,q_2,l_1,l_2\right)\in\mathbb{Z}_+^4 \tag{$\widetilde{R}3$}$$

Example 4. *The set* $\mathbb{R}_+^{\mathbb{Z}_+^2}$ *of all positive double sequences is regular.*

Example 5. *The set* $\widetilde{\mathcal{B}}$ *of all bounded sequences of* $\mathbb{R}_+^{\mathbb{Z}_+^2}$ *is regular.*

The following lemma, not difficult to prove, is needed in the formulation of the principal theorem of this paper.

Lemma 1. *Let* $\widetilde{\mathcal{R}}$ *be a regular subset of* $\mathbb{R}_+^{\mathbb{Z}_+^2}$*, then*

(*i*) *The subset* $\mathcal{R}^0:=\left\{N_{.,0}:N\in\widetilde{\mathcal{R}}\right\}$ *is regular in* $\mathbb{R}_+^{\mathbb{Z}_+}$.

(*ii*) *The subset* $\mathcal{R}_0:=\left\{N_{0,.}:N\in\widetilde{\mathcal{R}}\right\}$ *is regular in* $\mathbb{R}_+^{\mathbb{Z}_+}$.

3. THE ALGEBRA OF $\mathcal{R}-$REGULAR BOUNDED GENERALIZED FUNCTIONS

Let

$$\mathcal{S}^{*}\left(\Omega\right)=\left\{f\in\mathcal{C}^{\infty}\left(\Omega\right):\forall\alpha\in\mathbb{Z}_{+}^{n},\sup_{x\in\Omega}\left|\partial^{\alpha}f\left(x\right)\right|<\infty\right\},$$

and $\mathcal{R}$ be a regular subset of $\mathbb{R}_{+}^{\mathbb{Z}_{+}}$, if we define

$$\mathcal{E}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)=\left\{\left(u_{\epsilon}\right)_{\epsilon}\in\mathcal{S}^{*}\left(\Omega\right)^{I}:\exists N\in\mathcal{R},\forall\alpha\in\mathbb{Z}_{+}^{n},\sup_{x\in\Omega}\left|\partial^{\alpha}u_{\epsilon}\left(x\right)\right|=O\left(\epsilon^{-N_{\left|\alpha\right|}}\right),\epsilon\longrightarrow0\right\},$$

$$\mathcal{N}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)=\left\{\left(u_{\epsilon}\right)_{\epsilon}\in\mathcal{S}^{*}\left(\Omega\right)^{I}:\forall N\in\mathcal{R},\forall\alpha\in\mathbb{Z}_{+}^{n},\sup_{x\in\Omega}\left|\partial^{\alpha}u_{\epsilon}\left(x\right)\right|=O\left(\epsilon^{N_{\left|\alpha\right|}}\right),\epsilon\longrightarrow0\right\},$$

where $I=\left]0,1\right]$, then the properties of $\mathcal{E}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)$ and $\mathcal{N}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)$, easy to verify, are given by the following results.

Proposition 2. (i) *The space* $\mathcal{E}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)$ *is a subalgebra of* $\mathcal{S}^{*}\left(\Omega\right)^{I}$.
(ii) *The space* $\mathcal{N}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)$ *is an ideal of* $\mathcal{E}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)$.
(iii) *We have* $\mathcal{N}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)=\mathcal{N}_{\mathcal{S}^{*}}\left(\Omega\right)$, *where*

$$\mathcal{N}_{\mathcal{S}^{*}}\left(\Omega\right)=\left\{\left(u_{\epsilon}\right)_{\epsilon}\in\mathcal{S}^{*}\left(\Omega\right)^{I}:\forall m\in\mathbb{Z}_{+},\forall\alpha\in\mathbb{Z}_{+}^{n},\sup_{x\in\Omega}\left|\partial^{\alpha}u_{\epsilon}\left(x\right)\right|=O\left(\epsilon^{m}\right),\epsilon\longrightarrow0\right\}$$

We have also the null characterization of the ideal $\mathcal{N}_{\mathcal{S}^{*}}\left(\Omega\right)$ provided Ω is a box.

Definition 3. *An open subset* Ω *of* $\mathbb{R}^{n}$ *is said to be a box if*

$$\Omega=\mathbf{I}_{1}\times\mathbf{I}_{2}\times...\times\mathbf{I}_{n},$$

where each $\mathbf{I}_{i}$ *is a finite or infinite open interval in* $\mathbb{R}$.

Proposition 3. *Let* Ω *be a box, then an element* $\left(u_{\epsilon}\right)_{\epsilon}\in\mathcal{E}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)$ *belongs to* $\mathcal{N}_{\mathcal{S}^{*}}\left(\Omega\right)$ *if and only if the following condition is satisfied*

$$\forall m\in\mathbb{Z}_{+},\sup_{x\in\Omega}\left|u_{\epsilon}\left(x\right)\right|=O\left(\epsilon^{m}\right)\ ,\ \epsilon\longrightarrow0 \tag{1}$$

Proof. Suppose that $\left(u_{\epsilon}\right)_{\epsilon}\in\mathcal{E}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)$ satisfies (1). It suffices to show that $\left(\partial_{i}u_{\epsilon}\right)_{\epsilon}$ satisfies the $\mathcal{N}_{\mathcal{S}^{*}}\left(\Omega\right)$ estimates for all $i=1,..,n$. Suppose that u_{ϵ} is real valued, in the complex case, we shall carry out the following calculus separately on its real and imaginary part. Let $m\in\mathbb{Z}_{+}$, we have to show

$$\sup_{x\in\Omega}\left|\partial_{i}u_{\epsilon}\left(x\right)\right|=O\left(\epsilon^{m}\right),\ \epsilon\longrightarrow0$$

Since $\left(u_{\epsilon}\right)_{\epsilon}\in\mathcal{E}_{\mathcal{S}^{*}}^{\mathcal{R}}\left(\Omega\right)$, then

$$\exists N\in\mathcal{R},\sup_{x\in\Omega}\left|\partial_{i}^{2}u_{\epsilon}\left(x\right)\right|=O\left(\epsilon^{-N_{2}}\right),\ \epsilon\longrightarrow0 \tag{2}$$

Since $\left(u_{\epsilon}\right)_{\epsilon}$ satisfies (1), we have

$$\forall m\in\mathbb{Z}_{+},\sup_{x\in\Omega}\left|u_{\epsilon}\left(x\right)\right|=O\left(\epsilon^{N_{2}+2m}\right),\ \epsilon\longrightarrow0 \tag{3}$$

By Taylor's formula, we have

$$u_{\epsilon}\left(x+\epsilon^{N_{2}+m}e_{i}\right)=u_{\epsilon}\left(x\right)+\partial_{i}u_{\epsilon}\left(x\right)\epsilon^{N_{2}+m}+\frac{1}{2}\partial_{i}^{2}u_{\epsilon}\left(x+\theta\epsilon^{N_{2}+m}e_{i}\right)\epsilon^{2\left(N_{2}+m\right)},$$

where $\theta \in]0,1[$ and ϵ is sufficiently small as Ω is a box. It follows that

$$|\partial_i u_\epsilon(x)| \le \underbrace{\left|u_\epsilon\left(x+\epsilon^{N_2+m}e_i\right)\right|\epsilon^{-N_2-m}}_{(*)} + \underbrace{|u_\epsilon(x)|\,\epsilon^{-N_2-m}}_{(**)} + \underbrace{\epsilon^{N_2+m}\left|\partial_i^2 u_\epsilon\left(x+\theta\epsilon^{N_2+m}e_i\right)\right|}_{(***)}$$

From (3), we have $(*)$ and $(**)$ are of order $O(\epsilon^m)$, $\epsilon \longrightarrow 0$; and from (2), we have $(***)$ is of order $O(\epsilon^m)$, $\epsilon \longrightarrow 0$. $\square$

Definition 4. *Let $\mathcal{R}$ be a regular subset of $\mathbb{R}_+^{\mathbb{Z}_+}$, the algebra of $\mathcal{R}-$regular bounded generalized functions, denoted $\mathcal{G}^{\mathcal{R}}_{\mathcal{S}^*}(\Omega)$, is the quotient algebra*

$$\mathcal{G}^{\mathcal{R}}_{\mathcal{S}^*}(\Omega) = \frac{\mathcal{E}^{\mathcal{R}}_{\mathcal{S}^*}(\Omega)}{\mathcal{N}_{\mathcal{S}^*}(\Omega)} \tag{4}$$

Remark 1. *When $\mathcal{R}$ is the set of all positive sequences the algebra $\mathcal{G}^{\mathcal{R}}_{\mathcal{S}^*}(\Omega)$ is denoted by $\mathcal{G}_{L^\infty}(\Omega)$ in* [2] *and* [6] *as it is constructed on the differential algebra $D_{L^\infty}(\Omega)$ of Schwartz* [19]. *So it is more correct to write $\mathcal{G}^{\mathcal{R}}_{L^\infty}(\Omega)$ instead of $\mathcal{G}^{\mathcal{R}}_{\mathcal{S}^*}(\Omega)$. The null characterization of negligible elements of $\mathcal{G}_{L^\infty}(\Omega)$ in the case $\Omega = \mathbb{R}^n$ is given in* [9].

4. THE ALGEBRA OF $\mathcal{R}-$REGULAR ROUGHLY DECREASING GENERALIZED FUNCTIONS

Let

$$\mathcal{S}_*(\Omega) = \left\{ f \in \mathcal{C}^\infty(\Omega) : \forall \beta \in \mathbb{Z}_+^n, \sup_{x\in\Omega} \left|x^\beta f(x)\right| < \infty \right\},$$

and $\mathcal{R}$ be a regular subset of $\mathbb{R}_+^{\mathbb{Z}_+}$, if we define

$$\mathcal{E}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega) = \left\{ (u_\epsilon)_\epsilon \in \mathcal{S}_*(\Omega)^I : \exists N \in \mathcal{R}, \forall \beta \in \mathbb{Z}_+^n, \sup_{x\in\Omega} \left|x^\beta u_\epsilon(x)\right| = O\left(\epsilon^{-N_{|\beta|}}\right), \epsilon \longrightarrow 0 \right\},$$

$$\mathcal{N}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega) = \left\{ (u_\epsilon)_\epsilon \in \mathcal{S}_*(\Omega)^I : \forall N \in \mathcal{R}, \forall \beta \in \mathbb{Z}_+^n, \sup_{x\in\Omega} \left|x^\beta u_\epsilon(x)\right| = O\left(\epsilon^{N_{|\beta|}}\right), \epsilon \longrightarrow 0 \right\},$$

then the following properties of $\mathcal{E}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega)$ and $\mathcal{N}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega)$ are easy to verify.

Proposition 4. (i) *The space $\mathcal{E}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega)$ is a subalgebra of $\mathcal{S}_*(\Omega)^I$.*

(ii) *The space $\mathcal{N}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega)$ is an ideal of $\mathcal{E}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega)$.*

(iii) *We have $\mathcal{N}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega) = \mathcal{N}_{\mathcal{S}_*}(\Omega)$, where*

$$\mathcal{N}_{\mathcal{S}_*}(\Omega) = \left\{ (u_\epsilon)_\epsilon \in \mathcal{S}_*(\Omega)^I : \forall m \in \mathbb{Z}_+, \forall \beta \in \mathbb{Z}_+^n, \sup_{x\in\Omega} \left|x^\beta u_\epsilon(x)\right| = O(\epsilon^m), \epsilon \longrightarrow 0 \right\}$$

The following proposition characterizes $\mathcal{N}_{\mathcal{S}_*}(\Omega)$

Proposition 5. *Let $(u_\epsilon)_\epsilon \in \mathcal{E}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega)$, then $(u_\epsilon)_\epsilon \in \mathcal{N}_{\mathcal{S}_*}(\Omega)$ if and only if the following condition is satisfied*

$$\forall m \in \mathbb{Z}_+, \sup_{x\in\Omega} |u_\epsilon(x)| = O(\epsilon^m) \ , \ \epsilon \longrightarrow 0 \tag{5}$$

Proof. Suppose that $(u_\epsilon)_\epsilon \in \mathcal{E}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega)$ satisfies (5), since $(u_\epsilon)_\epsilon \in \mathcal{E}^{\mathcal{R}}_{\mathcal{S}_*}(\Omega)$, then $\exists N \in \mathcal{R}$, $\forall \beta \in \mathbb{Z}_+^n$,

$$\sup_{x\in\Omega} \left|x^{2\beta} u_\epsilon(x)\right| = O\left(\epsilon^{-N_{2|\beta|}}\right) \ , \ \epsilon \longrightarrow 0$$

From (5), we have

$$\forall m \in \mathbb{Z}_+, \sup_{x\in\Omega} |u_\epsilon(x)| = O\left(\epsilon^{2m+N_{2|\beta|}}\right), \ \epsilon \longrightarrow 0$$

Therefore $\forall x \in \Omega$,

$$\left|x^{\beta}u_{\epsilon}(x)\right|^{2} = \left|x^{2\beta}u_{\epsilon}(x)\right|\left|u_{\epsilon}(x)\right| = O\left(\epsilon^{2m}\right),\ \epsilon \longrightarrow 0,$$

hence

$$\forall m \in \mathbb{Z}_{+}, \left|x^{\beta}u_{\epsilon}(x)\right| = O\left(\epsilon^{m}\right),\ \epsilon \longrightarrow 0$$

□

Definition 5. *Let $\mathcal{R}$ be a regular subset of $\mathbb{R}_{+}^{\mathbb{Z}_{+}}$, the algebra of $\mathcal{R}-$regular roughly decreasing generalized functions, denoted $\mathcal{G}_{\mathcal{S}_{*}}^{\mathcal{R}}(\Omega)$, is the quotient algebra*

$$\mathcal{G}_{\mathcal{S}_{*}}^{\mathcal{R}}(\Omega) = \frac{\mathcal{E}_{\mathcal{S}_{*}}^{\mathcal{R}}(\Omega)}{\mathcal{N}_{\mathcal{S}_{*}}(\Omega)}$$

Remark 2. *The C^{∞} regularity in the definition of elements of $\mathcal{G}_{\mathcal{S}_{*}}^{\mathcal{R}}(\Omega)$ is not in fact needed in the proof of the principal results of this work.*

5. The algebra of $\mathcal{R}-$regular rapidly decreasing generalized functions

Let

$$\mathcal{S}(\Omega) = \left\{ f \in \mathcal{C}^{\infty}(\Omega) : \forall(\alpha,\beta) \in \mathbb{Z}_{+}^{2n}, \sup_{x\in\Omega}\left|x^{\beta}\partial^{\alpha}f(x)\right| < \infty \right\},$$

called the space of rapidly decreasing functions on Ω, and let $\widetilde{\mathcal{R}}$ be a regular subset of $\mathbb{R}_{+}^{\mathbb{Z}_{+}^{2}}$, if we define

$$\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega) = \left\{ (u_{\epsilon})_{\epsilon} \in \mathcal{S}(\Omega)^{I} : \exists N \in \widetilde{\mathcal{R}}, \forall(\alpha,\beta) \in \mathbb{Z}_{+}^{2n}, \sup_{x\in\Omega}\left|x^{\beta}\partial^{\alpha}u_{\epsilon}(x)\right| = O\left(\epsilon^{-N_{|\alpha|,|\beta|}}\right), \epsilon \longrightarrow 0 \right\},$$

$$\mathcal{N}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega) = \left\{ (u_{\epsilon})_{\epsilon} \in \mathcal{S}(\Omega)^{I} : \forall N \in \widetilde{\mathcal{R}}, \forall(\alpha,\beta) \in \mathbb{Z}_{+}^{2n}, \sup_{x\in\Omega}\left|x^{\beta}\partial^{\alpha}u_{\epsilon}(x)\right| = O\left(\epsilon^{N_{|\alpha|,|\beta|}}\right), \epsilon \longrightarrow 0 \right\},$$

then we have the following results.

Proposition 6. *We have the following assertions*

(i) The space $\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$ is a subalgebra of $\mathcal{S}(\Omega)^{I}$.

(ii) The space $\mathcal{N}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$ is an ideal of $\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$.

(iii) We have $\mathcal{N}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega) = \mathcal{N}_{\mathcal{S}}(\Omega)$, where

$$\mathcal{N}_{\mathcal{S}}(\Omega) = \left\{ (u_{\epsilon})_{\epsilon} \in \mathcal{S}(\Omega)^{I} : \forall m \in \mathbb{Z}_{+}, \forall(\alpha,\beta) \in \mathbb{Z}_{+}^{2n}, \sup_{x\in\Omega}\left|x^{\beta}\partial^{\alpha}u_{\epsilon}(x)\right| = O\left(\epsilon^{m}\right), \epsilon \longrightarrow 0 \right\}$$

Proof. The proof is not difficult, it follows from the properties of the set $\widetilde{\mathcal{R}}$. □

Definition 6. *Let $\widetilde{\mathcal{R}}$ be a regular subset of $\mathbb{R}_{+}^{\mathbb{Z}_{+}^{2}}$, the algebra of $\widetilde{\mathcal{R}}-$regular rapidly decreasing generalized functions on Ω, denoted by $\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$, is the quotient algebra*

$$\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega) = \frac{\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)}{\mathcal{N}_{\mathcal{S}}(\Omega)}$$

Example 6. *(i) For $\widetilde{\mathcal{R}} = \mathbb{R}_{+}^{\mathbb{Z}_{+}^{2}}$, we obtain the algebra $\mathcal{G}_{\mathcal{S}}(\Omega)$ of rapidly decreasing generalized functions on Ω, see* [13].

(ii) Pour $\widetilde{\mathcal{R}} = \widetilde{\mathcal{B}}$, we obtain the algebra $\mathcal{G}_{\mathcal{S}}^{\infty}(\Omega)$ of regular rapidly decreasing generalized functions on Ω, see [8].

6. CHARACTERIZATION OF $\mathcal{R}-$REGULAR RAPIDLY DECREASING GENERALIZED FUNCTIONS

Let us mention that the theorem of [4] can be extended to an open subset Ω of $\mathbb{R}^n$ provided Ω is a box.

Proposition 7. *If* Ω *is a box of* $\mathbb{R}^n$, *then*

$$\mathcal{S}(\Omega)=\mathcal{S}^*(\Omega)\cap\mathcal{S}_*(\Omega) \tag{6}$$

Proof. The proof is the same as in [4], noting that, in Taylor's expansion, the hypothesis that Ω is a box assures that (x_1+h,x') stays in Ω for all $(x_1,x')\in\Omega$ and $h>0$ sufficiently small. $\square$

The principal result of this section is an extension of (6) to the algebra of $\widetilde{\mathcal{R}}$-regular rapidly decreasing generalized functions. It is the first characterization of the algebra $\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$.

Theorem 8. *If* Ω *is a box, then*

$$\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)=\mathcal{G}_{\mathcal{S}_*}^{\mathcal{R}_0}(\Omega)\cap\mathcal{G}_{\mathcal{S}^*}^{\mathcal{R}^0}(\Omega)$$

Proof. We have to show that $\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)=\mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}(\Omega)\cap\mathcal{E}_{\mathcal{S}^*}^{\mathcal{R}^0}(\Omega)$ and $\mathcal{N}_{\mathcal{S}}(\Omega)=\mathcal{N}_{\mathcal{S}_*}(\Omega)\cap\mathcal{N}_{\mathcal{S}^*}(\Omega)$. The inclusions $\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)\subset\mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}(\Omega)\cap\mathcal{E}_{\mathcal{S}^*}^{\mathcal{R}^0}(\Omega)$ and $\mathcal{N}_{\mathcal{S}}(\Omega)\subset\mathcal{N}_{\mathcal{S}_*}(\Omega)\cap\mathcal{N}_{\mathcal{S}^*}(\Omega)$ are obvious. In order to show the reverse inclusions, first let $(u_\epsilon)_\epsilon\in\mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}(\Omega)\cap\mathcal{E}_{\mathcal{S}^*}^{\mathcal{R}^0}(\Omega)$, then $(u_\epsilon)_\epsilon\in\mathcal{S}^*(\Omega)^I\cap\mathcal{S}_*(\Omega)^I=\mathcal{S}(\Omega)^I$. In order to show that $(u_\epsilon)_\epsilon$ satisfies the estimate of $\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$, set $x=(x_1,x')\in\mathbf{I}_1\times(\mathbf{I}_2\times\mathbf{I}_3\times\ldots\times\mathbf{I}_n):=\Omega$ and consider in first the case $x_1>0$. For $h>0$ sufficiently small, a Taylor's expansion of u_ϵ with respect to x_1 gives

$$u_\epsilon(x_1+h,x')=u_\epsilon(x_1,x')+h\partial_1u_\epsilon(x_1,x')+\frac{h^2}{2}\partial_1^2u_\epsilon(\xi,x'), \tag{7}$$

for $\xi\in\,]x_1,x_1+h[$. The hypothesis $(u_\epsilon)_\epsilon\in\mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}(\Omega)\cap\mathcal{E}_{\mathcal{S}^*}^{\mathcal{R}^0}(\Omega)$ gives

$$\exists L\in\mathcal{R}_0,\forall k\in\mathbb{Z}_+,\sup_{x_1>0}(1+|x|)^k|u_\epsilon(x)|=O\left(\epsilon^{-L_k}\right),\epsilon\longrightarrow0$$

$$\exists M\in\mathcal{R}^0,\sup_{x_1>0}\left|\partial_1^2u_\epsilon(x)\right|=O\left(\epsilon^{-M_2}\right),\epsilon\longrightarrow0$$

We have

$$\sup_{x_1>0}(1+|x|)^k|u_\epsilon(x_1+h,x')|\leq\sup_{x_1>0}(1+|(x_1+h,x')|)^k|u_\epsilon(x_1+h,x')|=O\left(\epsilon^{-L_k}\right),\epsilon\longrightarrow0$$

It follows from (7)

$$|\partial_1u_\epsilon(x_1,x')|\leq\frac{1}{h}\left[|u_\epsilon(x_1+h,x')|+|u_\epsilon(x_1,x')|\right]+\frac{h}{2}\left|\partial_1^2u_\epsilon(\xi,x')\right|$$

Therefore

$$\sup_{x_1>0}(1+|x|)^k|\partial_1u_\epsilon(x)|=O\left(\epsilon^{-L_k-M_2}\right),\epsilon\longrightarrow0$$

From $\left(\widetilde{R}3\right)$ of definition 2 , there exists $N'\in\widetilde{\mathcal{R}}$ such that

$$L_k+M_2\leq N'_{2,k}\ ,$$

consequently

$$\forall\beta\in\mathbb{Z}_+^n,\sup_{x_1>0}\left|x^\beta\partial_1u_\epsilon(x)\right|\leq C\sup_{x_1>0}(1+|x|)^{|\beta|}|\partial_1u_\epsilon(x)|=O\left(\epsilon^{-N'_{2,|\beta|}}\right),\epsilon\longrightarrow0$$

If $x_1 < 0$, consider v_ϵ such that $v_\epsilon(x) = u_\epsilon(-x_1, x')$. We see that $(v_\epsilon)_\epsilon \in \mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}(\Omega) \cap \mathcal{E}_{\mathcal{S}^*}^{\mathcal{R}^0}(\Omega)$ and consequently the above arguments give the existence of $N" \in \widetilde{\mathcal{R}}$ such that

$$\sup_{x_1>0} \left| x^\beta \partial_1 v_\epsilon(x) \right| = \sup_{x_1<0} \left| x^\beta \partial_1 u_\epsilon(x) \right| = O\left(\epsilon^{-N"_{2,|\beta|}}\right), \epsilon \longrightarrow 0$$

Now from $\left(\widetilde{R}1\right)$ and $\left(\widetilde{R}2\right)$ of definition 2, there exists $N \in \widetilde{\mathcal{R}}$ such that

$$\max\left(N'_{2,|\beta|}, N"_{2,|\beta|}\right) \leq N_{1,|\beta|} \ ,$$

so

$$\sup_{x\in\Omega} \left| x^\beta \partial_1 u_\epsilon(x) \right| = O\left(\epsilon^{-N_{1,|\beta|}}\right), \epsilon \longrightarrow 0$$

Analogously we show

$$\exists N \in \widetilde{\mathcal{R}}; \forall \beta \in \mathbb{Z}_+^n, \sup_{x\in\Omega} \left| x^\beta \partial_i u_\epsilon(x) \right| = O\left(\epsilon^{-N_{1,|\beta|}}\right), i = 2, ..., n$$

Therefore, by induction, we obtain

$$\exists N \in \widetilde{\mathcal{R}}; \forall \alpha \in \mathbb{Z}_+^n, \forall \beta \in \mathbb{Z}_+^n, \sup_{x\in\Omega} \left| x^\beta \partial^\alpha u_\epsilon(x) \right| = O\left(\epsilon^{-N_{|\alpha|,|\beta|}}\right), \epsilon \longrightarrow 0,$$

i.e. $(u_\epsilon)_\epsilon \in \mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$.

Suppose now that $(u_\epsilon)_\epsilon \in \mathcal{N}_{\mathcal{S}_*}(\Omega) \cap \mathcal{N}_{\mathcal{S}^*}(\Omega)$, then

$$\forall m \in \mathbb{Z}_+, \forall k \in \mathbb{Z}_+, \sup_{x_1>0} (1+|x|)^k |u_\epsilon(x)| = O\left(\epsilon^{\frac{m}{2}}\right), \epsilon \longrightarrow 0$$

$$\forall m \in \mathbb{Z}_+; \sup_{x_1>0} \left| \partial_1^2 u_\epsilon(x) \right| = O\left(\epsilon^{\frac{m}{2}}\right), \epsilon \longrightarrow 0$$

We have

$$\sup_{x_1>0} (1+|x|)^k |u_\epsilon(x_1+h, x')| \leq \sup_{x_1>0} (1+|(x_1+h, x')|)^k |u_\epsilon(x_1+h, x')| = O\left(\epsilon^{\frac{m}{2}}\right), \epsilon \longrightarrow 0$$

It follows from (7)

$$\sup_{x_1>0} (1+|x|)^k |\partial_1 u_\epsilon(x)| = O(\epsilon^m), \epsilon \longrightarrow 0$$

Consequently

$$\forall m \in \mathbb{Z}_+, \forall \beta \in \mathbb{Z}_+^n, \sup_{x_1>0} \left| x^\beta \partial_1 u_\epsilon(x) \right|^2 \leq C_1 \sup_{x_1>0} (1+|x|)^{|\beta|} |\partial_1 u_\epsilon(x)| = O(\epsilon^m), \epsilon \longrightarrow 0$$

If $x_1 < 0$, consider v_ϵ such that $v_\epsilon(x) = u_\epsilon(-x_1, x')$ as above, then we obtain

$$\sup_{x_1>0} \left| x^\beta \partial_1 v_\epsilon(x) \right|^2 = \sup_{x_1<0} \left| x^\beta \partial_1 u_\epsilon(x) \right|^2 = O(\epsilon^m), \epsilon \longrightarrow 0$$

Therefore, by induction, we have

$$\forall m \in \mathbb{Z}_+, \forall \alpha \in \mathbb{Z}_+^n, \forall \beta \in \mathbb{Z}_+^n, \sup_{x\in\Omega} \left| x^\beta \partial^\alpha u_\epsilon(x) \right| = O(\epsilon^m), \epsilon \longrightarrow 0$$

Thus $\mathcal{N}_{\mathcal{S}_*}(\Omega) \cap \mathcal{N}_{\mathcal{S}^*}(\Omega) \subset \mathcal{N}_{\mathcal{S}}(\Omega)$ and consequently $\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega) = \mathcal{G}_{\mathcal{S}_*}^{\mathcal{R}_0}(\Omega) \cap \mathcal{G}_{\mathcal{S}^*}^{\mathcal{R}^0}(\Omega)$. □

The propositions 3 and 5 give the following result which characterizes the negligible elements of the algebra $\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$.

Corollary 9. *If Ω is a box, then an element $(u_\epsilon)_\epsilon \in \mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}(\Omega)$ is in $\mathcal{N}_{\mathcal{S}}(\Omega)$ if and only if the following condition is satisfied*

$$\forall m \in \mathbb{Z}_+, \sup_{x\in\Omega} |u_\epsilon(x)| = O(\epsilon^m), \epsilon \longrightarrow 0$$

7. CHARACTERIZATION OF $\mathcal{R}$−REGULAR RAPIDLY DECREASING GENERALIZED FUNCTIONS VIA FOURIER TRANSFORM

The direct Fourier transform of $u \in \mathcal{S}$, denoted $\widehat{u}$, is defined by

$$\widehat{u}(\xi) = (2\pi)^{-\frac{n}{2}} \int e^{-ix\xi} u(x)\, dx$$

Definition 7. *The Fourier transform of $u = [(u_\epsilon)_\epsilon] \in \mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}$, denoted by $\mathcal{F}_{\mathcal{S}}(u)$, is defined by*

$$\mathcal{F}_{\mathcal{S}}(u) = \widehat{u} = [(\widehat{u}_\epsilon)_\epsilon] \text{ in } \mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}$$

Remark 3. *The inverse Fourier transform of $u \in \mathcal{S}$, denoted $\widetilde{u}$, and the map $\mathcal{F}_{\mathcal{S}}^{-1}$ are defined as usually and in the same way.*

The following proposition gives one of the main results of the Fourier transform $\mathcal{F}_{\mathcal{S}}$ and is easy to prove.

Proposition 10. *The map*

$$\mathcal{F}_{\mathcal{S}} : \mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}} \longrightarrow \mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}$$

is an algebraic isomorphism.

Let

$$\widehat{\mathcal{S}^*} = \left\{ f \in \mathcal{C}^\infty : \forall \beta \in \mathbb{Z}_+^n, \sup_{\xi\in\mathbb{R}^n} \left|\xi^\beta \widehat{f}(\xi)\right| < \infty \right\},$$

and let $\widetilde{\mathcal{R}}$ be a regular subset of $\mathbb{R}_+^{\mathbb{Z}_+^2}$, if we define

$$\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0} = \left\{ (u_\epsilon)_\epsilon \in \widehat{\mathcal{S}^*}^I : \exists N \in \mathcal{R}^0, \forall \beta \in \mathbb{Z}_+^n, \sup_{\xi\in\mathbb{R}^n} \left|\xi^\beta \widehat{u_\epsilon}(\xi)\right| = O\left(\epsilon^{-N_{|\beta|}}\right), \epsilon \longrightarrow 0 \right\},$$

$$\mathcal{N}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0} = \left\{ (u_\epsilon)_\epsilon \in \widehat{\mathcal{S}^*}^I : \forall N \in \mathcal{R}^0, \forall \beta \in \mathbb{Z}_+^n, \sup_{\xi\in\mathbb{R}^n} \left|\xi^\beta \widehat{u_\epsilon}(\xi)\right| = O\left(\epsilon^{N_{|\beta|}}\right), \epsilon \longrightarrow 0 \right\},$$

then the following proposition is easy to prove.

Proposition 11. (i) *The space $\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$ is a subalgebra of $\widehat{\mathcal{S}^*}^I$.*

(ii) *The space $\mathcal{N}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$ is an ideal of $\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$.*

(iii) *The ideal $\mathcal{N}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0} = \mathcal{N}_{\widehat{\mathcal{S}^*}}$, where*

$$\mathcal{N}_{\widehat{\mathcal{S}^*}} := \left\{ (u_\epsilon)_\epsilon \in \widehat{\mathcal{S}^*}^I : \forall m \in \mathbb{Z}, \forall \beta \in \mathbb{Z}_+^n, \sup_{\xi\in\mathbb{R}^n} \left|\xi^\beta \widehat{u_\epsilon}(\xi)\right| = O(\epsilon^m), \epsilon \longrightarrow 0 \right\}$$

The following proposition characterizes $\mathcal{N}_{\widehat{\mathcal{S}^*}}$.

Proposition 12. *Let $(u_\epsilon)_\epsilon \in \mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$, then $(u_\epsilon)_\epsilon \in \mathcal{N}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$ if and only if the following condition is satisfied*

$$\forall m \in \mathbb{Z}_+, \sup_{\xi\in\mathbb{R}^n} |\widehat{u_\epsilon}(\xi)| = O(\epsilon^m) \ , \ \epsilon \longrightarrow 0 \tag{8}$$

Proof. The proof is similar to that of proposition 5. $\square$

Definition 8. *The algebra* $\mathcal{G}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$ *is defined as the quotient algebra*

$$\mathcal{G}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}=\frac{\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}}{\mathcal{N}_{\widehat{\mathcal{S}^*}}}$$

The next theorem is the second characterization of $\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}$.

Theorem 13. *We have*

$$\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}=\mathcal{G}_{\mathcal{S}_*}^{\mathcal{R}_0}\cap\mathcal{G}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$$

Proof. Let $(u_\epsilon)_\epsilon\in\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$, then $\exists C>0$ such that

$$\begin{aligned}\int\left|x^\beta\widehat{u_\epsilon}(x)\right|dx &\leq C\sup_{x\in\mathbb{R}^n}\left(1+|x|^2\right)^n\left|x^\beta\widehat{u_\epsilon}(x)\right|\\ &= O\left(\epsilon^{-N'_{|\beta|+2n}}\right),\ \epsilon\longrightarrow0\\ &= O\left(\epsilon^{-N_{|\beta|}}\right),\ \epsilon\longrightarrow0,\end{aligned}$$

for some $N\in\mathcal{R}^0$. The continuity of $\mathcal{F}$ from $\mathbb{L}^1$ to $\mathbb{L}^\infty$ gives

$$\left|\left|\partial^\beta u_\epsilon\right|\right|_{\mathbb{L}^\infty}=O\left(\epsilon^{-N_{|\beta|}}\right),\ \epsilon\longrightarrow0,$$

which shows that $(u_\epsilon)_\epsilon\in\mathcal{E}_{\mathcal{S}^*}^{\mathcal{R}^0}$ and therefore $\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}\subset\mathcal{E}_{\mathcal{S}^*}^{\mathcal{R}^0}$. Consequently $\mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}\cap\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}\subset\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}$. In order to show the inverse inclusion let us mention, at first, that from [4], we have

$$(u_\epsilon)_\epsilon\in\mathcal{S}^I\Longleftrightarrow(u_\epsilon)_\epsilon\in\mathcal{S}_*^I\bigcap\widehat{\mathcal{S}^*}^I$$

which implies in particular that $\mathcal{S}\subset\widehat{\mathcal{S}^*}$. On the other hand if $(u_\epsilon)_\epsilon\in\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}$, then

$$\begin{aligned}\int\left|\partial^\beta u_\epsilon(x)\right|dx &\leq C\sup_{x\in\mathbb{R}^n}\left(1+|x|^2\right)^n\left|\partial^\beta u_\epsilon(x)\right|\\ &= O\left(\epsilon^{-N'_{|\beta|,2n}}\right),\ \epsilon\longrightarrow0\\ &= O\left(\epsilon^{-N_{|\beta|,0}}\right),\ \epsilon\longrightarrow0,\end{aligned}$$

i.e.

$$\int\left|\partial^\beta u_\epsilon(x)\right|dx=O\left(\epsilon^{-N_{|\beta|}}\right),\ \epsilon\longrightarrow0,$$

for some $N\in\mathcal{R}^0$. The continuity of $\mathcal{F}$ from $\mathbb{L}^1$ to $\mathbb{L}^\infty$ gives

$$\left|\left|\xi^\beta\widehat{u_\epsilon}\right|\right|_{\mathbb{L}^\infty}=O\left(\epsilon^{-N_{|\beta|}}\right),\ \epsilon\longrightarrow0,$$

which shows that $(u_\epsilon)_\epsilon\in\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$ and consequently $(u_\epsilon)_\epsilon\in\mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}\cap\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$. Thus $\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}\subset\mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}\cap\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$, it follows that $\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}=\mathcal{E}_{\mathcal{S}_*}^{\mathcal{R}_0}\cap\mathcal{E}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$. A similar proof shows that $\mathcal{N}_{\mathcal{S}}=\mathcal{N}_{\mathcal{S}_*}\bigcap\mathcal{N}_{\widehat{\mathcal{S}^*}}$. Therefore $\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}=\mathcal{G}_{\mathcal{S}_*}^{\mathcal{R}_0}\cap\mathcal{G}_{\widehat{\mathcal{S}^*}}^{\mathcal{R}^0}$. $\square$

The following corollary gives a second characterization of the space $\mathcal{N}_{\mathcal{S}}$.

Corollary 14. *An element* $(u_\epsilon)_\epsilon\in\mathcal{E}_{\mathcal{S}}^{\widetilde{\mathcal{R}}}$ *is in* $\mathcal{N}_{\mathcal{S}}$ *if and only if the following condition is satisfied*

$$\forall m\in\mathbb{Z}_+,\ \sup_{\xi\in\mathbb{R}^n}\left|\widehat{u_\epsilon}(\xi)\right|=O\left(\epsilon^m\right),\epsilon\longrightarrow0 \tag{9}$$

8. Consequences

We know that when $\widetilde{\mathcal{R}} = \mathbb{R}_+^{\mathbb{Z}_+^2}$ we obtain $\mathcal{G}_{\mathcal{S}}^{\mathbb{R}_+^{\mathbb{Z}_+^2}} = \mathcal{G}_{\mathcal{S}}$. Theorem 8 gives the following corollary which is a characterization of the algebra of rapidly decreasing generalized functions.

Corollary 15. *We have*

$$\mathcal{G}_{\mathcal{S}} = \mathcal{G}_{\mathcal{S}^*} \cap \mathcal{G}_{\mathcal{S}_*} \ ,$$

where

$$\mathcal{G}_{\mathcal{S}^*} := \frac{\left\{ (u_\epsilon)_\epsilon \in \mathcal{S}^{*I} : \forall \alpha \in \mathbb{Z}_+^n, \exists m \in \mathbb{Z}_+, \sup\limits_{x \in \mathbb{R}^n} \left| \partial^\alpha u_\epsilon (x) \right| = O\left(\epsilon^{-m}\right), \epsilon \longrightarrow 0 \right\}}{\left\{ (u_\epsilon)_\epsilon \in \mathcal{S}^{*I} : \forall \alpha \in \mathbb{Z}_+^n, \forall m \in \mathbb{Z}_+, \sup\limits_{x \in \mathbb{R}^n} \left| \partial^\alpha u_\epsilon (x) \right| = O\left(\epsilon^{m}\right), \epsilon \longrightarrow 0 \right\}} \ ,$$

and

$$\mathcal{G}_{\mathcal{S}_*} := \frac{\left\{ (u_\epsilon)_\epsilon \in \mathcal{S}_*^{I} : \forall \beta \in \mathbb{Z}_+^n, \exists m \in \mathbb{Z}_+, \sup\limits_{x \in \mathbb{R}^n} \left| x^\beta u_\epsilon (x) \right| = O\left(\epsilon^{-m}\right), \epsilon \longrightarrow 0 \right\}}{\left\{ (u_\epsilon)_\epsilon \in \mathcal{S}_*^{I} : \forall \beta \in \mathbb{Z}_+^n, \forall m \in \mathbb{Z}_+, \sup\limits_{x \in \mathbb{R}^n} \left| x^\beta u_\epsilon (x) \right| = O\left(\epsilon^{m}\right), \epsilon \longrightarrow 0 \right\}}$$

We have also the following corollary which is an other characterization of the algebra $\mathcal{G}_{\mathcal{S}}$.

Corollary 16. *We have*

$$\mathcal{G}_{\mathcal{S}} = \mathcal{G}_{\mathcal{S}_*} \cap \mathcal{G}_{\widehat{\mathcal{S}^*}} \ ,$$

where

$$\mathcal{G}_{\widehat{\mathcal{S}^*}} := \frac{\left\{ (u_\epsilon)_\epsilon \in \widehat{\mathcal{S}^*}^{I} : \forall \beta \in \mathbb{Z}_+^n, \exists m \in \mathbb{Z}_+, \sup\limits_{\xi \in \mathbb{R}^n} \left| \xi^\beta \widehat{u_\epsilon} (\xi) \right| = O\left(\epsilon^{-m}\right), \epsilon \longrightarrow 0 \right\}}{\left\{ (u_\epsilon)_\epsilon \in \widehat{\mathcal{S}^*}^{I} : \forall \beta \in \mathbb{Z}_+^n, \forall m \in \mathbb{Z}_+, \sup\limits_{\xi \in \mathbb{R}^n} \left| \xi^\beta \widehat{u_\epsilon} (\xi) \right| = O\left(\epsilon^{m}\right), \epsilon \longrightarrow 0 \right\}}$$

We also know that when $\widetilde{\mathcal{R}} = \widetilde{\mathcal{B}}$ we obtain $\mathcal{G}_{\mathcal{S}}^{\widetilde{\mathcal{B}}} = \mathcal{G}_{\mathcal{S}}^{\infty}$. The next result, which is a corollary of theorem 8 for $\mathcal{R} = \widetilde{\mathcal{B}}$, gives a characterization of $\mathcal{G}_{\mathcal{S}}^{\infty}$.

Corollary 17. *We have*

$$\mathcal{G}_{\mathcal{S}}^{\infty} = \mathcal{G}_{\mathcal{S}^*}^{\infty} \cap \mathcal{G}_{\mathcal{S}_*}^{\infty} \ ,$$

where

$$\mathcal{G}_{\mathcal{S}^*}^{\infty} := \frac{\left\{ (u_\epsilon)_\epsilon \in \mathcal{S}^{*I} : \exists m \in \mathbb{Z}_+, \forall \alpha \in \mathbb{Z}_+^n, \sup\limits_{x \in \mathbb{R}^n} \left| \partial^\alpha u_\epsilon (x) \right| = O\left(\epsilon^{-m}\right), \epsilon \longrightarrow 0 \right\}}{\left\{ (u_\epsilon)_\epsilon \in \mathcal{S}^{*I} : \forall m \in \mathbb{Z}_+, \forall \alpha \in \mathbb{Z}_+^n, \sup\limits_{x \in \mathbb{R}^n} \left| \partial^\alpha u_\epsilon (x) \right| = O\left(\epsilon^{m}\right), \epsilon \longrightarrow 0 \right\}} \ ,$$

and

$$\mathcal{G}_{\mathcal{S}_*}^{\infty} := \frac{\left\{ (u_\epsilon)_\epsilon \in \mathcal{S}_*^{I} : \exists m \in \mathbb{Z}_+, \forall \beta \in \mathbb{Z}_+^n, \sup\limits_{x \in \mathbb{R}^n} \left| x^\beta u_\epsilon (x) \right| = O\left(\epsilon^{-m}\right), \epsilon \longrightarrow 0 \right\}}{\left\{ (u_\epsilon)_\epsilon \in \mathcal{S}_*^{I} : \forall m \in \mathbb{Z}_+, \forall \beta \in \mathbb{Z}_+^n, \sup\limits_{x \in \mathbb{R}^n} \left| x^\beta u_\epsilon (x) \right| = O\left(\epsilon^{m}\right), \epsilon \longrightarrow 0 \right\}}$$

We have also the following result obtained as a corollary of the theorem 13.

Corollary 18. *We have*

$$\mathcal{G}_{\mathcal{S}}^{\infty}=\mathcal{G}_{\mathcal{S}_*}^{\infty}\cap\mathcal{G}_{\widehat{\mathcal{S}^*}}^{\infty}$$

where

$$\mathcal{G}_{\widehat{\mathcal{S}^*}}^{\infty}:=\frac{\left\{(u_\epsilon)_\epsilon\in\widehat{\mathcal{S}^*}^{I}:\exists m\in\mathbb{Z}_+,\forall\beta\in\mathbb{Z}_+^n,\sup\limits_{\xi\in\mathbb{R}^n}\left|\xi^\beta\widehat{u_\epsilon}\left(\xi\right)\right|=O\left(\epsilon^{-m}\right),\epsilon\longrightarrow 0\right\}}{\left\{(u_\epsilon)_\epsilon\in\widehat{\mathcal{S}^*}^{I}:\forall m\in\mathbb{Z}_+,\forall\beta\in\mathbb{Z}_+^n,\sup\limits_{\xi\in\mathbb{R}^n}\left|\xi^\beta\widehat{u_\epsilon}\left(\xi\right)\right|=O\left(\epsilon^{m}\right),\epsilon\longrightarrow 0\right\}}$$

References

[1] J. Alvarez, H. Obiedat, Characterizations of the Schwartz space S and the Beurling-Björck space S_ω. Cubo 6, p. 167-183, (2004).

[2] H. Biagioni, M. Oberguggenberger, Generalized solutions to the Korteweg-de Vries and the generalized long-wave equations. SIAM J. Math Anal., 23(4), p. 923-940, (1992)

[3] C. Bouzar, K. Benmeriem, Ultraregular generalized functions. Oran-Essenia University, Preprint (2006).

[4] S.-Y. Chung, D. Kim et S. Lee. Characterization for Beurling-Bjorck space and Schwartz space. Proc. A.M.S. Vol. 125, N° 11, p. 3229-3234, (1997).

[5] J. F. Colombeau, Elementary introduction to new generalized functions, North Holland, (1985).

[6] J. F. Colombeau, A. Heibig, M. Oberguggenberger, Generalized solutions to PDEs of evolution type. Acta Appl. Math. 45, p.115–142. (1996)

[7] A. Delcroix, Fourier Transform of rapidly decreasing generalized functions. Application to microlocal regularity. J. Math. Anal. Appl., 327, p. 564-584, (2007).

[8] C. Garetto, Pseudo-differential operators in algebras of generalized functions and global hypoellipticity. Acta Appl. Math., 80(2):1, p. 123-174, (2004).

[9] C. Garetto, Topological structures in Colombeau algebras: investigation of the duals of $\mathcal{G}_c\left(\Omega\right),\mathcal{G}\left(\Omega\right)$ and $\mathcal{G}\left(\mathbb{R}^n\right)$. Monatsh. Math., 146(3), p. 203–226, (2005)

[10] S. Gindikin, L. Volevich. The Cauchy problem and related problems for convolution equations. Uspehi Mat. Nauk, vol. 27, no. 4, 65–143, (1972).

[11] K. Gröchenig, G. Zimmermann, Spaces of test functions via the STFT, J. Funct. Spaces Appl. 2, p. 25–53, (2004).

[12] I. M. Guelfand, G. E. Shilov. Generalized functions, vol. 2, Academic Press, (1967).

[13] M. Grosser, M Kunzinger, M. Oberguggenberger, R. Steinbauer . Geometric theory of generalized functions with applications to general relativity. Kluwer. (2001).

[14] L. Hörmander, Distributions theory and Fourier analysis. Springer, (1983).

[15] A. I. Kashpirovski, Equality of the spaces S_β^α and $S^\alpha\cap S_\beta$, Funct. Anal. Appl., 14, p. 129, (1980).

[16] M. Oberguggenberger, Regularity theory in Colombeau algebras. Bull. T. CXXXIII. Acad. Serbe Sci. Arts, Cl. Sci. Math. Nat. Sci. Math., 31, p. 147-162 , (2006).

[17] N. Ortner, P. Wagner, Applications of weighted $\mathcal{D}'_{\mathbb{L}^p}$-spaces to the convolution of distributions. Bull. Polish Acad. Sc. Math., Vol. 37, N^0 7-12, p. 579-595, (1989).

[18] Ya. V. Radyno, Sabra Ramdan, Ngo Fu Tkhan, The Fourier transform in an algebra of new generalized functions. Russian Acad. Sci. Dokl. Math. Vol. 46, N°3, p. 414-417 , (1993).

[19] L. Schwartz, Théorie des distributions. Herman, Paris, 2eme Ed., (1966).

Department of Mathematics, Oran-Essenia University, Oran, Algeria
E-mail address: `bouzar@univ-oran.dz; bouzarchikh@hotmail.com`

Centre Universitaire de Béchar, Algeria
E-mail address: `saidi_tb@yahoo.fr`